

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
Please do not report the images to the
Image Problem Mailbox.

Exhibit 2

TGF- β related genes in development

Nancy A Wall and Brigid LM Hogan

Vanderbilt University Medical School, Nashville, USA

Embryonic induction is the process by which signals from one cell population change the developmental fate of another. Polypeptides related to growth factors are one group of molecules mediating many inductive events. Recent data on the embryonic expression and function of signaling proteins related to transforming growth factor β , in both vertebrate and invertebrate systems, have shown that these molecules play important roles in both pattern formation and tissue specification during embryogenesis.

Current Opinion in Genetics and Development 1994, 4:517-522

Introduction

The transforming growth factor (TGF)- β superfamily is one of the largest groups of highly conserved intercellular signaling molecules regulating embryonic growth and differentiation. This superfamily, which currently includes at least 24 members, has several subgroups. The main groups are TGF- β s, *decapentaplegic*-Vg-related (DVR) proteins (including the bone morphogenetic proteins, or BMPs), and activins [1**]. Some members cannot be classified into a specific subgroup, however, and new ones are still being identified.

Over the past year, genetic, biochemical, and functional studies have supported the idea that TGF- β related proteins mediate inductive interactions during development. In some cases, these interactions involve a simple switch in the fate of the responding cells. In other systems, evidence suggests that TGF- β related proteins behave as morphogens, acting over relatively large distances in a graded fashion to differentially specify cell fate. This review includes data from studies in *Drosophila*, *Xenopus*, chick, and mouse, as well as *in vitro* studies with various cell lines.

Structural and biochemical analysis

TGF- β related proteins are synthesized as large pre-pro precursor molecules that are cleaved at an Arg-X-X-Arg site (where X can be any amino acid) to release a carboxy-terminal peptide of 110-140 amino acids, of which 7-9 are conserved cysteine residues. One of these cysteines is involved in intermolecular disulfide bonding to give a biologically active homo- or hetero-dimer [2,3].

Comparison of the crystal structures of three different growth factors, TGF- β 2, platelet-derived growth fac-

tor (PDGF)-BB, and nerve growth factor (NGF), has recently revealed a common structural motif involving two pairs of β -strands and an arrangement of conserved cysteine residues known as the 'cystine knot' [4]. Thus, although little overall sequence similarity is seen between these growth factors and they form dimers in different ways, they have a similar protomeric organization.

Recently, several new members of the DVR subgroup have been reported; these include nodal, glial-derived neurotrophic factor (GDNF), dorsalin-1, and two proteins which lack the conserved cysteine residue involved in dimerization, namely growth/differentiation factor-3/Vg-related 2 (GDF-3/Vgr-2) and GDF-9 [5,6,7**,8*,9**]. Fig. 1 shows the evolutionary relationships of DVR subgroup members. In some instances, the conservation of sequence means that vertebrate and invertebrate proteins can be functionally substituted; for example, two *Drosophila* proteins, DPP and 60A, induce bone when injected subcutaneously in rat [10]. Conversely, Padgett *et al.* [11] substituted the carboxy-terminal sequence of human BMP-4 with that of DPP to rescue *Drosophila decapentaplegic* (*dpp*) mutant embryos.

The biosynthesis of several TGF- β related proteins has now been studied in transfected cell lines. BMP-2, BMP-4, DPP, 60A, TGF- β 1, and DVR-6 are all processed similarly and, as predicted, dimers of cleaved mature region are secreted into the medium ([12-17]; NA Wall, unpublished data). The *in vivo* processing of TGF- β related proteins is much less well understood, however. BMP-1, though co-purified with several BMPs, does not resemble TGF- β , but rather has an astacin metalloproteinase domain and is closely related to the *Drosophila tolloid* gene product [18]. Null mutants of *tolloid* have a phenotype very similar to *dpp* mutants, and *tolloid* and *dpp* expression domains overlap [19]. Studies of *tolloid* mutants suggest that the protein is indirectly involved in proteolytically activating TGF-

Abbreviations

BMP—bone morphogenetic protein; *dpp*—*decapentaplegic*; DVR—*decapentaplegic*-Vg-related; GDF—growth/differentiation factor; GDNF—glial-derived neurotrophic factor; *hh*—*hedgehog*; NGF—nerve growth factor; PDGF—platelet-derived growth factor; TGF—transforming growth factor; *wg*—*wingless*.

8 Pattern formation and developmental mechanisms




Fig. 1. Phylogenetic relationships of decapentaplegic-Vg-related (DVR) proteins on the basis of comparison of carboxy-terminal sequences using the computer program PAUP 3.1 [59]. Similarity is indicated by the horizontal distance of any two sequences from a branch point. m—mouse; h—human; x—*Xenopus*; z—zebrafish; g—*Gallus* (chick); DPP—decapentaplegic; BMP—bone morphogenetic protein; OP—osteogenic protein; GDF—growth/differentiation factor; Vgr—Vg-related; GDNF—glial-derived neurotrophic factor.

β related proteins [20,21]. Recently, a *Xenopus* BMP-1 gene has been cloned and its expression shown to be tightly controlled during early development [22]. This raises the possibility that the activity of TGF- β related factors can be regulated *in vivo* at the level of protein processing.

New TGF- β related ligands continue to be identified, but relatively little is known about their receptors, which fall into two classes, type I and type II, both of which are families of transmembrane serine/threonine kinases. Whereas genes encoding receptors for TGF- β and activin have been cloned, only recently have candidate receptors for DVRs been reported. These include *daf-4*, a *Caenorhabditis elegans* type II receptor which can bind BMP-2 and BMP-4, and the *Drosophila saxophone* product, which appears to be a type I DPP receptor [23,24].

Genetic analysis of gene function

In the past year, several groups have published exciting new data on the function of *dpp* in *Drosophila*. Among other things, these studies raise the possibility that *dpp* regulates and is regulated by two other secreted growth factors, encoded by *wingless* (*wg*) and *hedgehog* (*hh*). As both *hh* and *wg*-type (Wnt) genes exist in vertebrates, work on *dpp* in *Drosophila* provides an important paradigm for the understanding of TGF- β related gene function in higher organisms.

The *dpp* gene is necessary for several different morphogenetic processes during *Drosophila* embryogenesis, including dorsal/ventral patterning of the embryo, establishment of the proximo-distal axis of appendages, eye development, and midgut morphogenesis [25–27]. In dorsal/ventral patterning, evidence suggests that the DPP protein acts as a morphogen regulating dorsal cell fate [28]. Whart n *et al.* [29**] have demonstrated that increasing the number of *dpp* genes in either mutant or

wild-type embryos results in the formation of an increasing number of dorsal type cells in a dose-dependent manner.

During midgut development, DPP is made in the visceral mesoderm and induces changes in gene expression in the adjacent endoderm, thus providing a model for understanding epithelial/mesenchymal interactions in higher organisms. Studies have shown that *dpp* transcription in the mesoderm is activated by the HOM gene *Ultrabithorax* (*Ubx*) and repressed by *Abdominal-A* (*Abd-A*) [30*] (Fig. 2a). In both cases, regulation is direct and involves multiple DNA-binding sites in a 5' upstream region of *dpp* [31*,32]. In the endoderm, DPP switches on the HOM gene *labial*, which has a DPP-response element in its 5' upstream region [33]. These studies clearly show that *dpp* acts during development both upstream and downstream of homeotic genes.

Studies on eye morphogenesis demonstrate yet another key role for *dpp* during *Drosophila* development [34**]. In this case, *dpp* is expressed in the morphogenetic furrow, a wave of organizing activity that moves across the eye imaginal disc (Fig. 2b). Transcription of *dpp* is regulated by the extracellular signaling molecule encoded by *hh*, and DPP induces morphogenesis in front of the furrow [35**]. In the leg disc, *dpp* is also regulated by *hh* [36**] and cooperates with *wg* in the regulation of the homeobox gene *aristaless*, which then acts as an organizer for proximal/distal patterning [37**] (Fig. 2c). In this case, at least three signaling molecules, *hh*, DPP, and *Wg*, coordinate to generate patterning in an epithelial sheet.

Drosophila is not the only system providing genetic clues to the biological importance of TGF- β related genes. Important contributions are now beginning to come from mice with mutations or targeted disruptions of genes in this family. The *short ear* (*se*) mutant has deletions or rearrangements in the gene for BMP-5 that are associated with loss of specific skeletal structures, indicating




Fig. 2. Schematic models for some of the interactions of *decapentaplegic* (*dpp*) with transcription factors and other signaling molecules during *Drosophila* development. (a) During visceral mesoderm/endoderm interactions in embryonic midgut development, *Ubx* directly and positively (+) regulates *dpp* transcription, whereas *Abd-A* inhibits it (-). The *DPP* protein elicits an intracellular signaling pathway in the endoderm that positively regulates *labial* transcription. *DPP* also indirectly maintains *Ubx* expression. Other interactions and regulatory factors are likely [30*,31*,33]. (b) During eye disc development, differentiated cells posterior to the morphogenetic furrow express *hedgehog* (*hh*). The diffusible *hh* signal regulates *dpp* expression in the cells of the morphogenetic furrow (anterior). *DPP* protein then mediates, directly or indirectly, morphogenesis anterior to the moving furrow [34**,35**]. (c) During leg disc development, cells expressing *engrailed* (*en*) in the posterior leg disc express *hh*, and *hh* diffuses anteriorly to adjacent cells to regulate *dpp* and *wingless* (*wg*) expression in dorsal and ventral regions, respectively. The interaction of *DPP* and *Wg* results in the focal transcriptional activation of the homeobox gene *aristaless* (*al*) which is thought to define a distal organizer [36**,37**].

a role for the gene product in skeletogenesis [38]. In the case of the mouse mutation 413-d, studies have shown that it results from a retroviral insertion in the TGF- β related gene *nodal*. During gastrulation, homozygous mutants are unable to form axial structures, including organized dorsal mesoderm and notochord [7**,39,40]. *In situ* hybridization studies have revealed that *nodal* RNA is expressed in a small group of cells around the node, but the precise role of the protein in axial patterning is not yet known.

Targeted disruptions of TGF- β 1, inhibin α , and activin/inhibin β B genes all result in aberrant phenotypes in homozygous mutant mice. Mice that are homozygous

null mutant for TGF- β 1 are viable for a few weeks before dying from a 'wasting' syndrome [41]. These results indicate that the embryo does not need to produce TGF- β 1 for prenatal development, although the mother may provide some TGF- β 1 protein in the uterus and milk. Mice deficient for inhibin α are also viable, but develop gonadal stromal tumors, suggesting that one function of the protein is to regulate or suppress stromal cell proliferation in this organ [42]. Targeted disruption of the related gene encoding activin/inhibin β B has also been reported [43]. Homozygous null mutants are viable, but females have reduced fertility, indicating a role for activin/inhibin β B in reproductive function. However, the fact that some viable offspring are born to these females

20 Pattern formation and developmental mechanisms

eliminates activin β B from playing a role in embryonic pattern formation.

Functional analysis

Besides genetic studies, various groups have used purified peptides and/or RNA misexpression to explore the function of TGF- β related proteins in a variety of *in vitro* assays. Overexpression of DVR-4/BMP-4 in *Xenopus* embryos causes induction of posterior and ventral types of mesoderm, whereas activin can induce dorsal types of mesoderm. In both cases, the proteins appear to act in part by regulating the expression of anterior-posterior-specific homeobox genes [44–48]. On the basis of the localization of its mRNA to the vegetal pole, Vg-1 was a prime candidate for an *in vivo* mesoderm inducer [49]. Initial experiments involving misexpression of Vg-1 mRNA did not show any effect on mesoderm formation, however. Although Vg-1 protein was produced, it was not processed to release the active, carboxy-terminal homodimer [50]. Recently, two groups have used constructs encoding chimeric proteins (BMP-2/Vg-1 and BMP-4/Vg-1) that circumvent the block on protein processing and have shown that mature Vg-1 does induce dorsal mesoderm [51*,52*]. Moreover, chimeric Vg-1 protein can rescue embryos ventralized by UV irradiation, suggesting that it acts by induction of the Nieuwkoop center. It now appears, therefore, that Vg-1 is an endogenous dorsal mesoderm inducer in *Xenopus* embryos, but is tightly regulated at a post-transcriptional level.

Two *in vitro* systems exploring epithelial/mesenchymal interactions in vertebrates have implicated a mediating role for BMP-4 and BMP-2. Vainio *et al.* [53*] have demonstrated that during tooth development, BMP-4 protein, which is initially expressed in presumptive dental epithelium, induces expression of several genes, including the endogenous BMP-4 gene, in adjacent mesenchyme. In the limb bud, where the apical ectodermal ridge influences the proliferation and differentiation of the underlying mesenchyme, Niswander and Martin [54*] have reported that BMP-2, which is transcribed in the apical ectodermal ridge, attenuates the proliferation of mesenchymal cells by fibroblast growth factor-4, a growth factor produced in the posterior apical ectodermal ridge.

In addition, BMP-2 modulates the differentiation of C3H10T1/2 cells, a mesenchymal stem cell line that can differentiate into myoblasts, chondrocytes, osteoblasts, and adipocytes. When treated with recombinant BMP-2, C3H10T1/2 cells differentiate into chondrocytes and osteoblasts at high protein concentrations and into adipocytes at low concentrations [55]. No myoblasts form in response to addition of BMP-2. In fact, Murray *et al.* [56] have reported that BMP inhibits expression of four myogenic determination genes, encoding myogenin, MyoD, herculin, and myf-5, in two myogenic cell lines. These results suggest that, *in vivo*, one factor regulating the differentiation of multipotent mesenchy-

mal stem cells into different mesodermal phenotypes is the local concentration of DVR protein to which they are exposed, either during embryogenesis or during processes such as wound healing and tissue repair.

Evidence for the patterning activity of TGF- β related proteins is also found in the developing nervous system. The *dorsalin-1* gene is normally transcribed in the dorsal roof plate of the chick embryonic spinal cord. When dorsalin-1 protein is added to cultured spinal cord explants, differentiation of neural crest cells is enhanced, whereas differentiation of ventral motor neurons is inhibited. It appears, therefore, that dorsalin-1 regulates the fate of dorsal cell types in the embryonic neural tube. A distantly related protein, GDNF, is capable of acting as a neurotrophic factor by specifically promoting the survival of midbrain dopaminergic neurons *in vitro*. Although several other TGF- β members are expressed in the nervous system at various times during embryogenesis, it is not yet known precisely how they function [15,57,58].

Conclusions

TGF- β related genes encode a large and growing family of structurally related secreted signaling molecules. Expression studies reveal that many of these genes are active in embryonic tissues undergoing inductive interactions and patterning. Genetic and functional studies support the idea that TGF- β related proteins mediate these events by acting upstream and downstream of transcription factors and other growth factors.

Future research will continue to examine the functional roles of these proteins during development. Targeted gene disruption in mice and interbreeding of different mutant lines will help to reveal which developmental processes require members of this superfamily family, and the production of purified proteins and antibodies will provide tools for elucidating the mechanisms involved. Also, identification of more receptors will facilitate our understanding of the signaling pathways activated by TGF- β related proteins. In fact, investigating these intracellular pathways appears to be one of the most important future directions to be taken in studying the role of TGF- β related genes during development. Given the high degree of sequence identity between members of the TGF- β superfamily, it will be interesting to determine the specificity or promiscuity of ligand-receptor interactions and to understand how the same ligand may elicit very different responses in different embryonic tissues. Finally, more data are needed about the relative roles of gene transcription, protein processing and degradation, and extracellular binding of mature protein in regulating the availability of ligand to receptors in responding tissues.

Acknowledgements

We would like to thank Dr CVE Wright for critical reading of the manuscript. Work in the authors' lab was supported in

part by grants HD28955 and CA48799. Brigid Hogan is an investigator of the Howard Hughes Medical Institute.

TGF- β related genes in development Wall and Hogan

521

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Kingsley DM: The TGF- β Superfamily: New Members, New Receptors, and New Genetic Tests of Function in Different Organisms. *Genes Dev* 1994, 8:133-146.
- A comprehensive and clearly written review covering all aspects of TGF- β with particular reference to genetic studies.
2. Daopin S, Piez KA, Ogawa Y, Davies DR: Crystal Structure of Transforming Growth Factor- β 2: an Unusual Fold for the Superfamily. *Science* 1992, 257:369-373.
3. Schlunegger MP, Gruter MC: An Unusual Feature Revealed by the Crystal Structure at 2.2 Å Resolution of Human Transforming Growth Factor- β 2. *Nature* 1992, 358:430-434.
4. McDonald NQ, Hendrickson WA: A Structural Superfamily of Growth Factors Containing a Cysteine Knot Motif. *Cell* 1992, 73:421-424.
5. Jones CM, Simon-Chazottes D, Cuenet JL, Hogan BL: Isolation of Vgr-2, a Novel Member of the Transforming Growth Factor- β -Related Gene Family. *Mol Endocrinol* 1992, 6:1961-1968.
6. McPherron AC, Lee S: GDF-3 and GDF-9: Two New Members of the Transforming Growth Factor- β Superfamily Containing a Novel Pattern of Cysteines. *J Biol Chem* 1993, 268:3444-3449.
7. Zhou X, Sasaki H, Lowe L, Hogan BLM, Kuehn MR: *nodal* is a Novel TGF- β -Like Gene Expressed in the Mouse Node during Gastrulation. *Nature* 1993, 361:543-547.
- This paper shows that *nodal* is related to TGF- β and is expressed in the node at a time and location coincident with dorsal mesoderm induction. The gene product may play a role in dorsal mesoderm induction and subsequent axial patterning.
8. Lin LH, Doherty DH, Lile JD, Bektesh S, Collins F: GDNF: A Glial Cell Line-Derived Neurotrophic Factor for Midbrain Dopaminergic Neurons. *Science* 1993, 260:1130-1132.
- This paper demonstrates a specific function for a TGF- β related gene in the nervous system. This raises the possibility that other members of this family may function as neurotrophic factors.
9. Basler K, Edlund T, Jessell TM, Yamada T: Control of Cell Pattern in the Neural Tube: Regulation of Cell Differentiation by *dorsalin-1* a Novel TGF- β Family Member. *Cell* 1993, 73:687-702.
- The cloning of the chick *dorsalin* gene is reported and *in situ* hybridization used to show the localization of its product to the dorsal embryonic spinal cord. *In vitro* culture assays were used to show that the protein enhances the differentiation of dorsal, but not ventral, cell types, suggesting that *in vivo*, *dorsalin* is involved in dorsal/ventral patterning of the neural tube.
10. Sampath TK, Rashka KE, Doctor JS, Tucker RF, Hoffmann FM: *Drosophila* Transforming Growth Factor Beta Superfamily Proteins Induce Endochondral Bone Formation in Mammals. *Proc Natl Acad Sci USA* 1993, 90:6004-6008.
11. Padgett RW, Wozney JM, Gelbart WM: Human BMP Sequences Can Confer Normal Dorsal-Ventral Patterning in the *Drosophila* Embryo. *Proc Natl Acad Sci USA* 1993, 90:2905-2909.
12. Panganiban FE, Rashka KE, Nietzel MD, Hoffmann FM: Biochemical Characterization of the *Drosophila* dpp Protein, a Member of the Transforming Growth Factor Beta Family of Growth Factors. *Mol Cell Biol* 1990, 10:2669-2677.
13. Miyazono K, Olofsson A, Colosetti P, Heldin C: A Role of the Latent TGF- β 1-Binding Protein in the Assembly and Secretion of TGF- β 1. *EMBO J* 1991, 10:1091-1101.
14. Doctor JS, Jackson D, Rashka KE, Visalli M, Hoffmann M: Sequence, Biochemical Characterization, and Developmental Expression of a New Member of the TGF- β Superfamily in *Drosophila melanogaster*. *Dev Biol* 1992, 151:491-505.
15. Wall NA, Blessing M, Wright CVE, Hogan BLM: Biosynthesis and *In Vivo* Localization of the Decapentaplegic-Vg-Related Protein, DVR-6 (Bone Morphogenetic Protein-6). *J Cell Biol* 1992, 120:493-502.
16. Israel DI, Nove J, Kerns KM, Moutsatsos IK, Kaufman RJ: Expression and Characterization of Bone Morphogenetic Protein-2 in Chinese Hamster Ovary Cells. *Growth Factors* 1992, 7:139-150.
17. Suzuki A, Nishimatsu S, Shoda A, Takebayashi K, Murakami K, Ueno N: Biochemical Properties of Amphibian Bone Morphogenetic Protein-4 Expressed in CHO Cells. *Biochem J* 1993, 291:413-417.
18. Wozney JM: Bone Morphogenetic Proteins. *Prog Growth Factor Res* 1990, 1:267-280.
19. Shimell MJ, Ferguson EL, Childs SR, O'Connor MB: The *Drosophila* Dorsal-Ventral Patterning Gene *tolloid* is Related to Human Bone Morphogenetic Protein 1. *Cell* 1991, 67:469-481.
20. Childs SR, O'Connor MB: Two Domains of the *tolloid* Protein Contribute to Its Unusual Genetic Interaction with *decapentaplegic*. *Dev Biol* 1994, 162:209-220.
21. Finelli AL, Bossie CA, Xie T, Padgett RW: Mutational Analysis of the *Drosophila* *tolloid* Gene, a Human BMP-1 Homolog. *Development* 1994, in press.
22. Meano M, Xue Y, Wood TI, Ong RC, Kung HF: Cloning and Expression of cDNA Encoding *Xenopus laevis* Bone Morphogenetic Protein-1 during Early Embryonic Development. *Gene* 1993, 134:257-261.
23. Estevez M, Attisano L, Wranna JL, Albert PS, Massague J, Riddle DL: The *daf-4* Gene Encodes a Bone Morphogenetic Protein Receptor Controlling *C. elegans* Dauer Larva Development. *Nature* 1993, 365:644-649.
24. Xie T, Finelli AL, Padgett RW: The *Drosophila* *saxophone* Gene: a Serine-Threonine Kinase Receptor of the TGF- β Superfamily. *Science* 1994, 263:1756-1759.
25. Spencer FA, Hoffmann FM, Gelbart WM: *Decapentaplegic* a Gene Complex Affecting Morphogenesis in *Drosophila melanogaster*. *Cell* 1982, 28:451-461.
26. Irish VF, Gelbart WM: The *decapentaplegic* Gene is Required for Dorsal-Ventral Patterning in the *Drosophila* Embryo. *Genes Dev* 1987, 1:868-879.
27. Panganiban CEF, Reuter R, Scott MP, Hoffmann FM: A *Drosophila* Growth Factor Homolog, *decapentaplegic*, Regulates Homeotic Gene Expression Within and Across Germ Layers during Midgut Morphogenesis. *Development* 1990, 110:1041-1050.
28. Ferguson EL, Anderson KV: Localized Enhancement and Repression of the Activity of the TGF- β Family Member, *decapentaplegic*, is Necessary for Dorsal-Ventral Pattern Formation in the *Drosophila* Embryo. *Development* 1992, 114:583-597.
29. Wharton KA, Ray RP, Gelbart WM: An Activity Gradient of *decapentaplegic* is Necessary for the Specification of Dorsal Pattern Elements in the *Drosophila* Embryo. *Development* 1993, 117:807-822.
- This paper shows that increasing numbers of *dpp* gene copies give increasingly dorsal-type tissue. This demonstrates that the dorsal/ventral character of embryonic tissue corresponds to high/low amounts of DPP and indicates that DPP can act as a morphogen.
30. Hursh DA, Padgett RW, Gelbart WM: Cross Regulation of *decapentaplegic* and *Ultrabithorax* Transcription in the Embryonic Visceral Mesoderm of *Drosophila*. *Development* 1993, 117:1211-1222.
- See [31].
31. Capovilla M, Brandt M, Botas J: Direct Regulation of *decapentaplegic* by *Ultrabithorax* and Its Role in *Drosophila* Midgut Morphogenesis. *Cell* 1994, 76:461-475.

522 Pattern formation and developmental mechanisms

This paper and [30*] show the direct regulation of a TGF- β related gene by a homeobox gene.

32. Masucci JD, Hoffman MF: Identification of Two Regions from the *Drosophila decapentaplegic* Gene Required for Embryonic Midgut Development and Larval Viability. *Dev Biol* 1993, 159:276-287.
33. Tremml C, Bienz M: Induction of *labial* Expression in the *Drosophila* Endoderm: Response Elements for *dpp* Signaling and for Autoregulation. *Development* 1992, 116:447-456.
34. Heberlein U, Wolff T, Rubin GM: The TGF- β Homolog *dpp* and the Segment Polarity Gene *hedgehog* Are Required for Propagation of a Morphogenetic Wave in the *Drosophila* Retina. *Cell* 1993, 75:913-926.
- This paper demonstrates that *hh*, which is expressed in cells posterior to the morphogenetic furrow, regulates expression of *dpp* within the furrow. DPP protein is required for progression of the furrow and, therefore, differentiation of the eye.
35. Ma C, Zhou Y, Beachy PA, Moses K: The Segment Polarity Gene *hedgehog* is Required for Progression of the Morphogenetic Furrow in the Developing *Drosophila* Eye. *Cell* 1993, 75:927-938.
- This paper demonstrates that morphogenetic furrow progression is halted by a temperature-sensitive allele of *hh*. The authors show that other genes involved in eye morphogenesis, *dpp* and *hairy*, are controlled by *hh*.
36. Basler K, Struhl G: Compartment Boundaries and the Control of *Drosophila* Limb Pattern by *hedgehog* Protein. *Nature* 1994, 368:208-214.
- Evidence is presented that, in the *Drosophila* wing and leg imaginal discs, the signaling protein *hedgehog* is produced by cells in the posterior compartment. In the leg disc, *hh* regulates the expression of DPP or Wg in a thin stripe of anterior cells adjacent to the compartment boundary. DPP then acts on cells on either side of the stripe to organize growth and patterning.
37. Campbell C, Weaver T, Tomlinson A: Axis Specification in the Developing *Drosophila* Appendage: the Role of *wingless*, *decapentaplegic*, and the Homeobox Gene *aristaeless*. *Cell* 1993, 74:1113-1123.
- The authors demonstrate that *wg* and *dpp* act together to induce *aristaeless* (*al*) expression. The region of *al* expression becomes the focal point from which a proximal/distal axis extends. Ectopic expression of *wg* or *al* in an area of high *dpp* expression results in a duplication of the proximal/distal axis.
38. Kingsley DM, Bland AE, Grubbs JM, Marker PC, Russell LB, Copeland NG, Jenkins NA: The Mouse Short Ear Skeletal Morphogenesis Locus is Associated with Defects in a Bone Morphogenetic Member of the TGF Beta Superfamily. *Cell* 1992, 71:399-410.
39. Conlon FL, Barth KS, Robertson EJ: A Novel Retrovirally Induced Embryonic Lethal Mutation in the Mouse: Assessment of the Developmental Fate of Embryonic Stem Cells Homozygous for the 413.1 Proviral Integration. *Development* 1991, 111:969-981.
40. Iannaccone PM, Zhou X, Khokha M, Boucher D, Kuehn MR: Insertional Mutation of a Gene Involved in Growth Regulation of the Early Mouse Embryo. *Dev Dynam* 1992, 194:198-208.
41. Shull MM, Ormsby I, Kier AB, Pawlowski S, Diebold RJ, Yin M, Allen R, Sidman C, Proetzel G, Calvin D, et al.: Targeted Disruption of the Mouse Transforming Growth Factor- β Gene Results in Multifocal Inflammatory Disease. *Nature* 1992, 359:693-699.
42. Matzuk MM, Finegold MJ, Su JJ, Hsueh AJW, Bradley A: α -Inhibin is a Tumour-Suppressor Gene with Gonadal Specificity in Mice. *Nature* 1992, 360:313-319.
43. Vassalli A, Matzuk MM, Gardner HAR, Lee K, Jaenisch R: Activin/Inhibin β 8 Subunit Gene Disruption Leads to Defects in Eyelid Development and Female Reproduction. *Genes Dev* 1994, 8:414-427.
44. Jones CM, Lyons KM, Lapan PM, Wright CVE, Hogan BLM: DVR-4 (Bone Morphogenetic Protein-4) as a Posterior-Ventralizing Factor in *Xenopus* Mesoderm Induction. *Development* 1992, 115:639-647.
45. Dale L, Howes G, Price BMJ, Smith JC: Bone Morphogenetic Protein 4: a Ventralizing Factor in Early *Xenopus* Development. *Development* 1992, 115:573-585.
46. Kimelman D, Christian JL, Moon RT: Synergistic Principles of Development: Overlapping Patterning Systems in *Xenopus* Mesoderm Induction. *Development* 1992, 116:1-9.
47. Sive HL: The Frog Prince-ss: a Molecular Formula for Dorsal-Ventral Patterning. *Genes Dev* 1993, 7:1-12.
48. Slack JMW: Inducing Factors in *Xenopus* Early Embryos. *Curr Biol* 1994, 4:116-126.
49. Weeks DL, Melton DA: A Maternal mRNA Localized to the Vegetal Hemisphere in *Xenopus* Eggs Codes for a Growth Factor Related to TGF- β . *Cell* 1987, 51:861-867.
50. Tannahill D, Melton DA: Localized Synthesis of the Vg-1 Protein during Early *Xenopus* Development. *Development* 1989, 106:775-785.
51. Thomsen GH, Melton DA: Processed Vg1 Protein is an Axial Mesoderm Inducer in *Xenopus*. *Cell* 1993, 74:433-441.
- This paper demonstrates that active Vg-1 peptide is a potent dorsal mesoderm inducer, by using a chimeric protein that circumvents the block on proteolytic processing of this protein. This indicates that protein processing is a critical point of regulation during development.
52. Dale L, Matthews G, Colman A: Secretion and Mesoderm-Inducing Activity of the TGF-Beta-Related Domain of *Xenopus* Vg1. *EMBO J* 1993, 12:4471-4480.
- A chimeric protein containing the proregion and proteolytic cleavage site of BMP-4 linked to the mature region of *Xenopus* Vg-1 is used to finally demonstrate mesoderm-inducing activity of Vg-1.
53. Vainio S, Karavanova I, Jowett A, Thesleff I: Identification of BMP-4 as a Signal Mediating Secondary Induction between Epithelial and Mesenchymal Tissues during Early Tooth Development. *Cell* 1993, 75:45-58.
- In vitro* culture was used to show that BMP-4 can induce differentiation of dental epithelium. This demonstrates that BMP-4 is autoinductive and shows a direct role for BMP-4 in epithelial/mesenchymal interaction.
54. Niswander L, Martin GR: FGF-4 and BMP-2 Have Opposite Effects on Limb Growth. *Nature* 1993, 361:68-71.
- This study employed an *in vitro* culture system to demonstrate fibroblast growth factor (FGF)-4 and BMP-2 effects on limb growth. FGF-4 is stimulatory and BMP-2 is inhibitory. This demonstrates that growth factors provide both stimulatory and inhibitory activity in regulation of limb growth.
55. Wang EA, Israel DI, Kelly S, Luxenberg DP: Bone Morphogenetic Protein-2 Causes Commitment and Differentiation in C3H10T1/2 and 3T3 Cells. *Growth Factors* 1993, 9:57-71.
56. Murray SS, Murray JB, Clackson CA, Urist MR: Bone Morphogenetic Protein Inhibits Differentiation and Affects Expression of Helix-Loop-Helix Regulatory Molecules in Myoblastic Cells. *J Cell Biochem* 1993, 53:51-60.
57. Flanders KC, Ludecke C, Engels S, Cissel DS, Roberts A, Kondaiah P, Lafyatis R, Sporn MB, Unsicker K: Localization and Actions of Transforming Growth Factor- β s in the Embryonic Nervous System. *Development* 1991, 113:183-191.
58. Lee SJ: Expression of Growth/Differentiation Factor 1 in the Nervous System: Conservation of a Bicistronic Structure. *Proc Natl Acad Sci USA* 1991, 88:4250-4254.
59. Swofford DL, Olver SH: Inferring Evolutionary Trees from Gene Frequency Data under the Principle of Maximum Parsimony. *Sys Zool* 1987, 36:293-325.

BLM Hogan, Howard Hughes Medical Institute, Vanderbilt University School of Medicine C-2310, Medical Center North, Nashville, Tennessee 37209, USA.

NA Wall, Department of Cell Biology, Vanderbilt University School of Medicine C-2310, Medical Center North, Nashville, Tennessee 37209, USA.